

A GIS Model for Predicting the Real-Time Risk of Arboviral Transmission in FL - Jon Day

- a) System
 - i) Arbovirus - SLE
 - ii) Primary vector - *Cx nigripalpus*
 - iii) Amplification
 - (1) Mosquitoes
 - (2) Birds
 - iv) Vector
 - (1) Extreme drought will shut down breeding
 - (2) Extreme flooding will as well
 - (3) Floodwater Culex
 - (4) Oviposition sites are in low-lying areas where periodic flooding occurs
- b) Setup
 - i) 589 MWTD data recording stations across peninsular FL
 - ii) Can measure rise and fall of ground water as well as other parameters
 - iii) See a lot of variability in water table depth
- c) Historic data
 - i) 1977 - SLE outbreak
 - (1) Water table showed a typical dry season dry down
 - (2) Mid May - June initial wetting
 - (3) Secondary dry down
 - (4) Wet season increase
 - ii) 1990 - SLE outbreak
 - (1) Similar profile
 - (2) Defined dry and wet seasons
 - iii) Particular interest in the initial wetting period
- d) Data Modeling
 - i) Model water table depth
 - ii) Look at how typical the water table data are for epidemic transmission
 - iii) Turn data into a risk map
 - (1) Take daily values at each MWTD site
 - (2) Graph how closely data match risk model
 - (3) Map data to risk map
- e) Pinellas County Case Study - 2005
 - i) 3 of 5 major SLE epidemics began in this county
 - ii) 2005
 - (1) 18 cases
 - (2) Onset dates - 1 June to 22 Aug
 - (3) 6/3/05 risk map showed high risk in Pinellas County area
 - (4) 8/12/05 risk map showed decreasing risk in area
 - iii) County Mosquito Control tracked virus in mosquito pools and birds
 - iv) WNV
 - (1) Mapped cases against the risk graphs
 - (2) Human cases occurred when data matched model
 - (3) Mosquito populations

- (a) High in year prior to epidemic
- (b) Populations skewed to older mosquitoes during epidemic year
- (4) Sentinels showed positives to match human cases
- f) What is the effect of human populations density?